

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 693 498 A1

B1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:

24.01.1996 Bulletin 1996/04

(51) Int. Cl.<sup>6</sup>: C07H 19/06, C07D 473/00,  
A61K 31/70, A61K 31/52

(21) Application number: 95114526.7

(22) Date of filing: 11.04.1990

(84) Designated Contracting States:

AT BE CH DE DK ES FR GB GR IT LI LU NL SE

(30) Priority: 17.04.1989 GB 8908646

20.09.1989 GB 8921266

(62) Application number of the earlier application in  
accordance with Art. 76 EPC: 90303904.8

(71) Applicant: SCOTIA HOLDINGS PLC  
Guildford Surrey GU1 1BA (GB)

(72) Inventors:

- Horrobin, David Frederick,  
c/o Scotia Pharm. Ltd.  
Guildford, Surrey GU1 1BA (GB)

- Winther, Michael David,  
c/o Scotia Pharm. Ltd.  
Guildford, Surrey GU1 1BA (GB)
- Stewart, John Charles Marshall,  
c/o Scotia Pharm.  
Guildford, Surrey GU1 1BA (GB)

(74) Representative: Sturt, Clifford Mark et al  
London WC1R 4JH (GB)

Remarks:

This application was filed on 15 - 09 - 1995 as a  
divisional application to the application mentioned  
under INID code 60.

(54) **Antiviral compounds**

(57) Anti-virals wherein linoleoyl or alpha-linolenyl or  
an unsaturated long chain fatty acyl group of the n-7 or  
n-9 series is borne directly on a hydroxy or amino group  
of the sugar/sugar analogue or heterocyclic moiety of a  
nucleoside or nucleoside analogue.

EP 0 693 498 A1


# Description

## FIELD OF INVENTION


The invention relates to anti-viral agents; they are fatty acid nucleosides and nucleoside analogue derivatives, and corresponding compounds with the fatty acid groups carried on an amino rather than hydroxy function of the nucleoside or nucleoside analogue.

## GENERAL


A number of compounds have been shown to exert anti-viral actions both in vitro and in vivo, in particular various purine and pyrimidine derivatives with a glycoside or glycoside analogue structure. Examples include:-




ACV (acyclovir;  
9-(2-hydroxyethoxymethyl)  
guanine)




DHPG  
(gancyclovir; 9-(1,3-  
dihydroxy-s-  
propoxymethyl)guanine)




AZT  
(azidothymidine,  
zidovudine)




D4T  
(dideoxy-didehydro-  
thymidine)



ddC  
(dideoxycytidine)



AHT  
(anhydrothymidine)



ddA  
(dideoxyadenosine)

In use, all these compounds, which are analogues of natural nucleosides, have limited ranges of action and even in vitro they affect only a limited range of viruses. Their mode of action is primarily to interfere with one or other of the stages of viral replication and assembly, an action referred to as replication inhibition herein. They do not inactivate the virus when it is present extra-cellularly and they have little or no effect on the transmission of active virus from cell to cell.

There are agents of a different kind that are able to inactivate viruses and inhibit their transmission, especially those viruses which are enveloped. These agents include long chain unsaturated fatty acids and their derivatives such as esters, salts, amides and glycerides. The effects of these agents are referred to as virus inactivation herein, but there may also be limited inhibitory effects on replication. It is thought that a major component of the virus inactivation is simply the result of the dramatic disruption of the protecting virus envelope.

Compounds in which nucleosides and nucleoside analogues are derivatised with fatty acids have been previously described as possessing anti-viral activity. Hamamura et al (J. Med. Chem. 19(5), 1976, 654-62; J. Med. Chem. 19(5), 1976, 663-67 and J. Med. Chem. 19(5), 1976, 667-73) disclose arabinofuranosyl cytosine derivatives with saturated and mono-unsaturated long chain fatty acids, while Ishida (JP-A-76-32573) describes derivatives of this nucleoside with linoleic acid. EP-A-0082668 discloses deoxyuridine analogues derivatised with saturated or unsaturated C<sub>1</sub> to C<sub>18</sub> fatty acids. EP-A-0085424 describes nucleoside analogues, such as gancyclovir, with linoleic and alpha-linolenic acids.

## THE INVENTION

The invention has several distinct but overlapping and inter-related aspects described primarily in relation to AZT, AHT, and acyclovir derivatives but extending equally to ddC, ddA, D4T and gancyclovir derivatives.

At its broadest the invention provides new anti-viral compounds, being long chain, unsaturated fatty acid ester or amide derivatives of nucleosides and nucleoside analogues, the fatty acid acyl groups being borne directly on a hydroxy or amino group of the sugar/sugar-analogue or heterocyclic moiety of the nucleoside or nucleoside analogue. The fatty acids are unsaturated long chain fatty acyl groups of the natural n-7 and n-9 series (C:16 upwards) and the parent acids of the n-6 and n-3 series, linoleic acid and alpha-linolenic acid. Particular examples of the nucleoside analogues are AZT, AHT and ACV.

The invention in the present aspect also provides:-

(i) A method of enhancing transport of anti-virals across lipid barriers in the body, and especially from the gut into the lymph system, or into cells from the extra-cellular fluid, or across the blood brain barrier, or percutaneously in topical preparations, or of inhibiting glucuronidation or other metabolic modifications reducing drug efficacy, in the treatment of viral infections, wherein said anti-viral is used in the form of a compound as above.

(ii) A method of preparation of a medicament for such purposes, wherein a compound as above is prepared as said medicament alone or in a pharmaceutically acceptable diluent or carrier.

In a second aspect the invention provides compounds being derivatives of pentose nucleosides, or of analogues thereof having an acyclic group in place of the pentose which acyclic group has a hydroxy substituted carbon at the position corresponding to the 5' position of the pentose, said derivatives being esters at the 5' position of the nucleoside or at the corresponding hydroxy substituted carbon of the analogue as the case may be, formed with the fatty acids referred to above, namely unsaturated long chain fatty acids of the n-9 and n-7 series (C:16 upwards), linoleic acid and alpha-linolenic acid.

Especially in the present aspect the invention provides AZT or AHT, or ACV, or ddA, ddC or D4T in the form of such compounds.

The invention in this aspect further provides:-

(i) A method of treatment of viral infections, with enhancement or inhibition as above but without loss of replication inhibiting activity, wherein said anti-viral is used in the form of a compound that is such a 5' or corresponding ester derivative.

(ii) A method of preparation of a medicament for such purposes, wherein such a compound is prepared as said medicament alone or in a pharmaceutically acceptable diluent or carrier.

In a third aspect the invention provides AZT and AHT, or ddA, ddC or D4T in the form of an ester derivative of the fatty acids referred to above, namely the unsaturated long chain fatty acids of the n-9 and n-7 series (C:16 upwards), linoleic acid and alpha-linolenic acid, active against herpes or other virus infection.

The invention in the present aspect further provides:-

(i) A method of treatment of herpes virus infection, particularly herpes simplex and herpes zoster or other sensitive viral infection, wherein there is administered to a sufferer therefrom an effective amount of such an AZT or AHT or ddA, ddC or D4T derivative.

(ii) Preparation of a medicament for the treatment of herpes virus infection, particularly herpes simplex and herpes zoster or other sensitive viral infection, wherein such an AZT or AHT or ddA, ddC or D4T derivative is prepared as said medicament, alone or in a pharmaceutically acceptable diluent or carrier.

In a fourth aspect the invention provides acyclovir or gancyclovir in the form of a virucidal ester or amide of a fatty acid as referred to above, namely the unsaturated long chain fatty acids of the n-9 and n-7 series (C:16 upwards), linoleic acid and alpha-linolenic acid.


In this aspect the invention further provides:-

- (i) A method of treatment of virus infection wherein there is administered to a sufferer therefrom an effective amount of such an acyclovir or gancyclovir derivative.
- (ii) Preparation of a medicament for the treatment of virus infection, wherein such an acyclovir or gancyclovir derivative is prepared as said medicament alone or in a pharmaceutically acceptable diluent or carrier.
- (iii) Such method, or preparation of medicament, wherein the infection is herpes simplex or herpes zoster or other sensitive viral infection.

Specific aspects of the invention may also be states as lying in:-

- (i) Particular derivatives of particular nucleoside analogue anti-virals, namely esters of AZT and AHT with long chain unsaturated fatty acids of the n-9 and n-7 series (C:16 upwards), linoleic acid and alpha-linolenic acid set out herein. These are believed to be new compounds, but the invention is more specifically in their application against herpes viruses, including herpes zoster and herpes simplex, where AZT and AHT are inactive. In this aspect therefore the invention lies in treating herpes infections with these esters, or in the preparation for such treatment of medicaments containing the esters.
- (ii) Particular derivatives of a particular nucleoside analogue anti-viral; namely esters or amides of acyclovir with long chain unsaturated fatty acids of the n-9 and n-7 series (C:16 upwards), linoleic acid and alpha-linolenic acid set out herein.

The formula of LA-acyclovir ester is, for example:



AK-6

where the pentose ring of AZT or AHT is represented by equivalents to the 1', 4' and 5'-carbon atoms of the ring. LA-acyclovir amide carries the linoleoyl group on the amino substituent of the purine nucleus, and di-LA-acyclovir at both positions.

These are again believed to be new compounds, but the invention also lies in treating virus infections, especially herpes virus infections with them, and in the preparation for use in such treatment of medicaments containing the esters.

## DISCUSSION

The fatty acid nucleoside derivatives have dramatically different physico-chemical properties from the parent nucleoside compounds. The compounds of the present invention are lipophilic with high absolute solubilities in chloroform and a large chloroform-water partition co-efficient (see Tables 3 and 4 herein).

The fatty acid nucleoside derivatives have one or more of the following advantages that are a consequence of new physical and biological properties:-

1. Combined virucidal and replication inhibition activities.
2. Broadening activity of replication inhibition to different viruses.

3. Enhanced penetration through and into the skin and into cells (topical formulations).
4. Improved pharmacokinetics (particularly in oral and i.v. formulations). These advantages are discussed in greater detail below.

#### 5 1. Combined virucidal and replication inhibition activities

The observation that two distinct modes of action against viruses can be combined in a single molecule are shown by most of the compounds of the invention is unexpected. Although free long chain unsaturated fatty acids are virucidal to most enveloped viruses, triglycerides are not, so clearly some chemical linkages interfere with this type of activity, LA in triglyceride format for example fails to retain the virucidal activity of the fatty acid component.

The acyclovir ester with LA (AK6) is also lacking in this activity, but AKs 2,4 and 11 are all virucidal.

The second unexpected property of the compounds is that they retain the replication inhibiting activity of the parent nucleoside when tested in in vitro assays for virus replication. This is surprising as many of the derivatives are 5'-hydroxy esters. For AZT and ACV it has been shown that the intra-cellular biologically active form is triphosphorylated at the 5'-hydroxy position. AZT and ACV, if tested lacking the phosphate groups, are inactive against viral polymerases.

There are two possible explanations as to the mode of action of the 5'-substituted compounds. The most likely is that the fatty acid group, having served its purpose in enhancing drug penetration through membranes, is rapidly cleaved off from the nucleoside by any of a variety of enzymes present in the cell cytoplasm. The de-acylated compound could then be a substrate for the phosphorylation needed to generate the active compound.

Less likely is the possibility that the fatty acid group in some way substitutes for the phosphorylation in the fatty acid-nucleoside, resulting in a compound which is active without cleavage and phosphorylation.

Results from anti-viral screening work are given later herein, and what stands out most clearly is that "AK 11" (the di-substituted ACV compound) is very effective both at inactivating virus and inhibiting replication.

#### 25 2. Broadening activity against viruses

A further unexpected observation is that, for example, LA-AZT gains the property of inhibiting herpes virus replication, a property not found with either LA or AZT individually (see Results section *infra*). The precise basis for this broadening of activity of an anti-retroviral agent to include herpes viruses is unknown but could be due to the enhanced permeability through cell membranes. Since multiple concomitant viral infections frequently occur it is of potential clinical benefit to treat them with a single active compound.

#### 3. Enhanced penetration through skin

Topical treatments for herpes virus infections were the first formulations to be developed for this virus. However, all such formulations produced to date have been of limited value for two reasons: firstly because the anti-viral agents are not lipophilic they do not easily penetrate the skin and therefore do not always achieve therapeutic concentrations at the site of virus replication. For this reason agents known as "enhancers" (e.g. dimethyl sulphoxide and propylene glycol) have been added to improve percutaneous absorption. However, most enhancers cause local toxicity and irritation with prolonged use. Thus, lipophilic derivatives of anti-virals that retain all of their potency in inhibiting virus replication are desirable.

The second reason for the disappointing results obtained with known topical treatments is that the virus replication occurs early in the natural history of infection, at the prodromal stage, usually before lesions are apparent. Thus any agent that acts only on the viral polymerase will be unable to halt further development of the lesion. However, an agent with virucidal properties, capable of disrupting the envelope structure of intact virus, may be expected to have superior efficacy in halting lesion development, reducing the chance of spreading infections virus to new locations and possibly reducing the subsequent load of latent virus that will lead to subsequent infections.

The compounds of the present invention, particularly AKs 11 and 13 are virucidal and are lipophilic so their penetration through the lipid matrix of the stratum corneum and also their penetration into cells will be efficient. These compounds will be present in the oil phase of oil-in-water or water-in-oil emulsions which simplifies the manufacture and application of such products.

#### 4. Improved pharmacokinetics

The metabolism and pharmacokinetics of fatty acid nucleosides are dramatically altered compared to the parent nucleosides. Some examples of how this will enhance the therapeutic efficacy are given.

a. Drug absorption

The lipid solubility of a drug is fundamental to absorption. For polar compounds (those which dissolve more easily in water than lipid) such as the parent nucleosides, crossing a non-porous, lipid membrane by diffusion is difficult and hence this process becomes rate limiting. Lipid soluble agents pass more easily through the barrier membranes of the gastro-intestinal tract which will increase their oral bioavailability. They will also tend to be absorbed at least in part by the lymphatic system. This latter route has the considerable advantage of avoiding the liver whereas water soluble agents are absorbed via the portal vein and so must go through the liver before reaching the general circulation. The "first pass metabolism" encountered during transport through the liver is a principal route of drug modification leading to excretion of inactive metabolites.

It is known that saturated fatty acids can enhance the transport of drugs across membranes but this has been believed efficient only with the shorter chain molecules of 6 - 12 carbons in length (for example see "Drug Delivery and Targeting Systems", 30.11.90. to 1.12.89. London, Conference documentation IBC Technical Services Ltd. Article by Professor Y. W. Chien). From this data it would have been expected that the longer chain unsaturated fatty acids of the present invention would be less effective, not active as they are.

b. Blood brain barrier

Many viruses (such as HIV and certain herpes viruses) are able to spread to the brain and cause infections leading to encephalitis or dementia. A critical factor in treatment is then the ability of a drug to pass through the blood brain barrier. It has been repeatedly observed that hydrophilic drugs do not readily pass the blood brain barrier and so only low levels can be detected in cerebrospinal fluid. Lipophilic drugs will more easily reach therapeutic levels in the brain. This is especially the case for the compounds of the present invention when given by the i.v. route.

c. Altered metabolism

Fifteen to twenty percent of zidovudine (AZT) is excreted unchanged into the urine and seventy five percent is metabolised by hepatic glucuronidation to form 3'-azido-2'-3'-dideoxy-5'-glucuronyl thymidine (an inert metabolite). Since derivatives such as AK2 (LA-AZT) have an ester linkage at this 5' position these will not be metabolised by the same pathway. This will increase the effective blood concentration and allow a lower dose of drug to achieve the same therapeutic benefit.

The combination of higher bioavailability and reduced metabolism will result in less variation in pharmacokinetics between individuals and so lead to lower doses and reduced incidence of toxic side effects.

There are further advantages to the production of amide derivatives of the anti-viral agents ddC and ddA. For both drugs the combined ester and amide linkages will increase lipophilicity and even if one of the fatty acids were to be removed by enzymic metabolism the remaining molecule will still be lipophilic and retain the advantages described.

For amide derivatives of ddA there is a further specific advantage in that this linkage will prevent the conversion of ddA to dId (dideoxyinosine) following deamination by adenosine deaminase which occurs within minutes of exposure to human plasma. If the fatty acid-ddA derivative can enter an HIV infected cell it can be converted to ddA and subsequently to the active triphosphate form. In contrast dId must eventually undergo further enzymic conversions to form ddA triphosphate (the active metabolite within the cell).

Furthermore, in order to be fully effective, most anti-viral agents must be present in both intra-cellular and extra-cellular compartments. Since the cell membrane is lipid rich, lipophilic compounds are more readily able to enter cells than are hydrophilic ones. Thus the drugs specified in the present invention are more likely to achieve effective intra-cellular concentrations than are their parent compounds.

## SUITABLE FATTY ACIDS AND SOURCES

Suitable unsaturated fatty acids include the natural n-9 and n-7 series fatty acids (C:16 upwards), linoleic acid and alpha-linolenic acid. The members of the n-6, n-3, n-7 and n-9 series of fatty acids up to C:22 are included in Table 1



below:

TABLE 1

	n-7 Diet or bodily synthesis	n-9 Diet or bodily synthesis	n-6 Diet	n-3 Diet
	16:0 (palmitic)	18:0 (stearic)		
delta 9-desaturase	↓	↓	↓	↓
	16:1 n-7 (palmitoleic)	18:1 n-9 (oleic)	18:2 n-6 (linoleic)	18:3 n-3 (alpha- linolenic)
delta 6-desaturase	↓	↓	↓	↓
	16:2 n-7	18:2 n-9	18:3 n-6 (gamma- linolenic)	18:4 n-3
	↓	↓	↓	↓
	18:2 n-7	20:2 n-9	20:3 n-6 (dihomo-gamma- linolenic)	20:4 n-3
delta 5-desaturase	↓	↓	↓	↓
	18:3 n-7	20:3 n-9	20:4 n-6 (arachidonic)	20:5 n-3
	↓	↓	↓	↓
	20:3 n-7	22:3 n-9	22:4 n-6 (adrenic)	22:5 n-3
delta 4-desaturase	↓	↓	↓	↓
	20:4 n-7	22:4 n-9	22:5 n-6	22:6 n-3

In the body the n-6 and n-3 series are dietary requirements, at least linoleic acid (LA) and alpha-linolenic acid (ALA) therefore being required as essential fatty acids. The n-7 and n-9 series are available endogenously, from palmitic acid (16:0) and stearic acid (18:0) respectively, by the action of delta-9 desaturase and the successive action of the other desaturases. The palmitic and stearic acids may be from dietary sources or bodily synthesis.

The position of the double bonds under the n-7, n-9 etc. nomenclature is counted from the non-carboxyl or omega end, with the unsaturations present in the chain as  $(-CH=CH-CH_2)_M$ , where M = 1 to 6, as represented in the terminology

18:1, 22:3, etc. Illustrations are:-

TABLE 2

n-6 series essential fatty acids		
18:2		delta-9,12 (linoleic acid)
18:3		delta-6,9,12 (gamma-linolenic acid)
20:3		delta-8,11,14 (dihomo-gamma-linolenic acid)
20:4		delta-5,8,11,14 (arachidonic acid)
22:4		delta-7,10,13,16 (adrenic acid)
22:5		delta-4,6,10,13,16
n-3 series essential fatty acids		
18:3		delta-9,12,15 (alpha-linolenic acid)
18:4		delta-6,9,12,15
20:4		delta-8,11,14,17
20:5		delta-5,8,11,14,17
22:5		delta-7,10,13,16,19
22:6		delta-4,7,10,13,16,19

The acids are systematically named as derivatives of the corresponding hexadecanoic, octadecanoic, eicosanoic or docosanoic acids, e.g. delta-9,12-octadecadienoic acid or delta 4,7,10,13,16,19-docosahexanoic acid, but numerical designations such as, correspondingly, 18:2 n-6 or 22:6 n-3 are convenient. Initials, for example, AA for arachidonic acid and more systematically DHA for 22:6 n-3 (docosahexanoic acid) or EPA for 20:5 n-3 (eicosapentaenoic acid), are also used but as to the latter kind do not serve when n-3 and n-6 acids of the same chain length and degree of unsaturation exist, as for example with the 22:5 acids. Trivial names in more or less common use in the n-6 series are as shown. Of the n-3 series only 18:3 n-3 has a commonly used trivial name, alpha-linolenic acid. It was characterised earlier than gamma-linolenic acid and reference in the literature simply to linolenic acid, especially in the earlier literature, is to the alpha-acid.

Fatty acids for use in the invention may be derived from natural sources, known in themselves and no part of the invention, or from chemical synthesis. For example oils derived from the seeds of certain plants such as *Oenothera biennis*, *Borago officinalis* and members of the *Ribes* family are often rich in 18:3 n-6. The oils from certain algae such as *Spirulina spp* and certain fungi such as *Mortierella spp* and *Rhizopus spp* may also contain 18:3 n-6. The 20:4 n-6 acid is found in egg yolks, in the lipids of certain fungi and algae, and in fish and other marine oils, especially those from warm waters. The 18:3 n-3 acid is widely distributed in plant oils, while 18:4 n-3 is found in the seed oils from members of the *Ribes* family. The 20:5 n-3 and 22:6 n-3 acids are often abundantly present in marine oils. Natural sources of 22:4 n-6 and 22:5 n-6 include adrenal glands (22:5 n-6) and kidneys (22:4 n-6) obtained from slaughter houses.

The n-9 fatty acid oleic acid (18:1 n-9) can be obtained from a large variety of plant oils, notably olive oil, and palmitoleic (16:1 n-7) can be found in most plant oils also.

## DOSAGES

It will be understood that the absolute quantity of active materials present in any dosage unit should not exceed that appropriate to the rate and manner of administration to be employed but on the other hand should also desirably be adequate to allow the desired rate of administration to be achieved by a small number of doses. The rate of administration will moreover depend on the precise pharmacological action desired.

The amounts of the compounds to be administered are 1 mg to 20 g per day in divided doses, preferably 10 mg to 4 g per day and more preferably 50 mg to 1 g per day and dosage unit forms are conveniently made up with these amounts or sub-multiples thereof. The compounds may be administered, orally, topically, enterally, parenterally (intravenously, intravascularly, intraperitoneally, intramuscularly or subcutaneously) or by any other appropriate route all per se known and using per se known presentations, though as already discussed herein there are advantages in particular modes of administration. Thus compounds may be administered as tablets, e.g. enteric-coated tablets to prevent destruction in the stomach, hard or soft gelatin capsules, syrups, emulsions, solutions, or any other form appropriate for oral or

parenteral administration; or in any form appropriate for topical administration such as ointments, creams, shampoos, lotions; or as pessaries and other formulations for rectal or vaginal application, or sticks. When applied topically the products may be prepared in concentrations of the active compounds of from 0.001 to 20%, preferably 0.1 to 10%, and especially preferably 1 to 5% by weight. Such a preparation is desirably used to give, at most, the daily dosages above and desirably dosages at the lower end of the ranges for example 20 mg/day.

## SYNTHESES


### Example 1. LA-AZT ("AK 2")

To a stirred solution of AZT (0.723g, 2.70 mmol, as a TLC-homogeneous gum) in anhydrous pyridine (10 ml), was added LA acid chloride (1.300 g, 4.36 mmol). The reaction was stirred under dry nitrogen at room temperature for a period of six hours. After this period, TLC (thin layer chromatography) indicated complete loss of starting material and the pyridine was removed in vacuo, evaporation temperature being maintained below 45°C. The resulting crude syrup was dissolved in chloroform: petrol (2:1, v:v) and was introduced onto a column of silica gel (30 g) pre-equilibrated in chloroform: petrol (2:1). Elution with chloroform: petrol (2:1) removed the unwanted high r.f. products. AK-2 was obtained as a colourless homogenous syrup by elution with chloroform: petrol (4:1). Yield 54% (0.770 g, 1.458 mmol).

### Example 2. LA-AHT ("AK 4")

Anhydrothymidine (0.643 g, 2.86 mmol, as a dry white powder) was dissolved in anhydrous pyridine (10 ml) under an atmosphere of dry nitrogen, and to this stirred solution at room temperature was added LA acid chloride (1.20 g, 4.00 mmol). Stirring was continued overnight after which period, TLC indicated complete loss of the starting material. The pyridine was evaporated in vacuo (below 45°C) and the residual syrup was dissolved in chloroform and then applied to a column of silica gel (30 g). Elution with chloroform: petrol (4:1, v:v) followed by chloroform and then chloroform: ethanol (50:1) gave the required derivative AK-4 as a colourless TLC-homogeneous gum. Yield, 55% (0.7653 g, 1.573 mmol).

### Example 3. O-LA-ACV. ("AK 6")



LA-OLEOYL ACID CHLORIDE

ACYCLOVIR


AK-6

Acyclovir (ACV, 0.5 g, 2.22 mmol, used as a dry white powder) was suspended in anhydrous pyridine (10 ml) and to the mixture stirred under dry nitrogen was added LA acid chloride (0.77 g, 2.58 mmol). The reaction mixture was stirred overnight at room temperature after which period TLC indicated conversion to the required product. The pyridine was removed in vacuo and the residual syrup was dissolved in chloroform. This was introduced into a column of silica gel (30 g) pre-equilibrated in chloroform: ethanol (30 : 1, v:v). Elution with chloroform followed by chloroform: ethanol (30:1), (20:1) and finally (15:1), gave the required product, AK-6, as a TLC homogeneous semi-solid. Yield, 45% (0.487 g, 1.0 mmol).

### Example 4. Di-LA-ACV. ("AK 11")

Acyclovir (ACV, 0.5 g, 2.22 mmol, used as a dry white powder) was suspended in anhydrous pyridine (10 ml). To the stirred mixture under dry nitrogen was added LA acid chloride (1.5 g, 5.1 mmol) and 4-di-methyl-amino pyridine (0.061 g, 0.5 mmol). The reaction was stirred overnight at room temperature after which period TLC indicated conversion to the required product. The pyridine was removed in vacuo and the residual syrup was dissolved in chloroform. This was introduced onto a column of silica gel (30 g) pre-equilibrated in chloroform. Elution with chloroform: petrol (3:1, v:v) followed by chloroform, gave the required product AK-11, as a TLC homogeneous syrup. Yield 60% (1.001 g, 1.332 mmol).

Example 5. n-LA-ACV ("AK 13")




AK-11

AK-13

A sample of AK-11 (0.786 g, 1.100 mmol) was dissolved in anhydrous methanol (10 ml) and anhydrous dioxane (2 ml). (The dioxane helps to solubilise the starting material). The solution was then cooled to 0°C and 0.30 ml of NaOMe solution (from Aldrich, 30% wt) was added. Stirring was maintained for fifteen minutes at 0°C after which period TLC indicated complete loss of the starting material and the formation of a slightly lower r.f. product. Dowex ion exchange resin (20 ml, in H<sup>+</sup> form) was added and stirring was maintained at room temperature for a further ten minutes. The Dowex was filtered and washed with methanol and the combined filtrate and washings were then evaporated in vacuo to afford the crude product as a pale yellow gum. The gum was dissolved in a minimum of chloroform and applied to a column of silica gel (5 g). Elution with chloroform and then chloroform : methanol (20:1) gave the required product as a white solid. The yield of AK 13 was 61% (0.326 g, 0.67 mmol).

Example 6. LA-D4T ("AK 15")



LA ACID CHLORIDE

D-4T

AK-15

A solution of dideoxy-didehydro thymidine (0.380 g, 1.727 mmol) in anhydrous pyridine (10 ml) was cooled to 0°C and to this solution was added LA acid chloride (1.033 g, 3.45 mmol). The solution was stirred overnight under an atmosphere of dry nitrogen after which period TLC indicated complete loss of the starting material. The pyridine was removed in vacuo and the residual syrup was dissolved in chloroform : petrol (2:1). This was then introduced onto a column of silica gel (20 g) pre-equilibrated in chloroform : petrol (2:1). Elution with chloroform : petrol (2:1), (3:1) and finally chloroform gave the required product as a TLC homogeneous syrup. Yield 53% (0.442 g, 0.92 mmol).

The following two synthesis examples are alternative routes for the AZT, AHT and ACV derivatives and are given in outline.

Example 7. LA - 3'azido-3'deoxythymidine. LA-AZT ("AK 2") (Alternative Method)

AZT (approx. 1 mmol) is dissolved in anhydrous pyridine (approx. 3 ml) and to this solution is added LA acid chloride (approx. 1 mmol). The solution is stirred at 0°C for one hour and then for period, the reaction mixture is evaporated to dryness to afford a syrup which is dissolved in dichloromethane (30 ml) and then washed with saturated sodium bicarbonate solution (20 ml). The organic layer is departed and evaporated in vacuo. The crude orange syrup is introduced onto the top of a 10 g column of silica gel. Elution with chloroform, followed by 20:1, 20:1, 10:1, CHCl<sub>3</sub> : MeOH mixtures enabled the separation of the required product.

The only difference with the other fatty acids is to use the corresponding acid chloride.

Example 8. LA-Anhydrothymidine. LA-AHT ("AK 4") (Alternative Method)

Procedure: As for LA-AZT except that chromatography requires 10% MeOH. As with LA-AZT, the only difference with other fatty acids is to use the corresponding acid chloride.

Example 9. Di-LA-ACV ("AK 11") (Alternative Method)

Acyclovir (approx. 3 mmol) is suspended in a stirred solution of pyridine (30 ml) and linoleoyl chloride (approx. 8 mmol) is slowly added to the pyridine. Acyclovir, which itself is insoluble in pyridine, gradually solubilises as linoleoyl esters are formed. The solution is stirred overnight and then poured into 200 ml of saturated sodium bicarbonate solution. The sodium bicarbonate solution is extracted twice with 150 ml chloroform. The chloroform fraction is dried with anhydrous sodium sulphate and then evaporated to dryness under vacuum. The resulting gum is applied to a silica gel column (200 mesh) in hexane and chromatographed with hexane/chloroform in increasing chloroform proportions, to give after evaporation di-linoleoyl acyclovir.

The only difference with the other fatty acids is to use the corresponding acid chloride, for example that of ALA.

The following synthesis examples are given in abbreviated form using the same methods but with different starting bases. They are written out as for LA derivatives but ALA derivatives and n-9 and n-7 series fatty acid derivatives are all, as with Examples 1-9, made simply by using the acid chlorides of the desired acids instead of those of LA expressly set out.

Example 10. o-LA-ddC "NL 1"

The procedure for this synthesis corresponds to that for AK2 but with use of ddC; a conventional protecting group is first added to the amino position of the base to restrict subsequent reaction with the linoleoyl acid chloride to the 5'-hydroxy group. After formation of the ester linkage the protecting group is removed leaving the title compound ("o" indicates attachment of the fatty acyl group at an oxygen atom, that of the 5'-hydroxy group).

Example 11. n-LA-ddC "NL 2"

The procedure for this synthesis again corresponds to that for AK2; no amino protecting group being used however. Since the amino position on the pyrimidine ring is more reactive than the hydroxy position the first acyl compound to form will be the n-LA-ddC (amide linkage, "n" indicating attachment of the fatty acyl group at a nitrogen atom, that of the heterocycle amino group).

Example 12. di-LA-ddC "NL 3"

The procedure for this synthesis is as above but carried to completion. Both reactive sites (hydroxyl and amine) react with the linoleoyl acid chloride.

Example 13. o-LA-ddA "NL 4"

The procedure for this synthesis is the same as that for NL1 except that ddA is used as the substrate base.

Example 14. n-LA-ddA "NL 5"

The procedure for this synthesis is the same as that for NL2 except that ddA is used as the substrate base.

Example 15. di-LA-ddA "NL 6"

The preparation of this material is the same as that for NL3 except that ddA is used as the substrate base.

Example 16. n-LA-DHPG "NL 7"

The preparation of this material is the same as AK13 except that gancyclovir is used in place of acyclovir. Initial reaction conditions result in both ester and amide linkages being formed as in the preparation of NL8, but ester bonds are subsequently broken by hydrolysis in alkaline conditions.

Example 17. tri-LA-DHPG "NL 8" (two ester linkages, one amide).

The preparation of this material corresponds to the preparation of AK11.

Example 18. o-di-LA-DHPG "NL 9"

The preparation of this material corresponds to the preparation of AK6.

#### EXAMPLES OF FORMULATIONS

The following examples of formulations illustrate the therapeutic application of the invention, in formulations which as to the mode of formulation per se are conventional as follows:-

##### a. Syrup of suspension

active ingredient	0.25 g
sorbitol solution	1.5 g
glycerol	0.005 g
dispersible cellulose	0.005 g
sodium benzoate	0.010 ml
water	5 litre

Mix the sorbitol and glycerol with part of the water. Dissolve the sodium benzoate in water and add to the bulk, then add and disperse the cellulose and active ingredient. Make up to volume.

##### b. Suppository

	mg/suppository
active ingredient	250
hard fat, BP (Witepsol H15-Dynamit Nobel (trade mark))	1770
	<u>2020</u>

One-fifth of the Witepsol H15 is melted at 45°C and the active ingredient added to the molten base with mixing. The remaining Witepsol H15 is added and stirred to a homogenous mix. The entire suspension is passed through a 250 µm stainless steel screen, cooled to 38°C to 40°C and filled into plastic moulds.

Administration by suppository gives access to the bloodstream without passage through the portal vein to the liver, where glucuronidation and other processes causing loss of active drug primarily occur.

c. Pessaries

	mg/pessary
active ingredient	250
anhydrous dextrose	380
potato starch	363
magnesium stearate	7
	<u>1000</u>

The above ingredients are mixed directly and pessaries prepared by direct compression of the resulting mixture.

Pessaries like the formulations at d), e) and j) are an example of topical formulations, showing the advantageous percutaneous absorption properties given by the fat-compatible nature of the derivatives of the invention.

d. Cream

	weight (g)
active ingredient	5.0
glycerol	2.0
cetostearyl alcohol	6.8
sodium lauryl sulphate	0.8
white soft paraffin	12.5
liquid paraffin	5.0
chlorocresol	0.1
purified water	100.00

The active compounds are dissolved in a mixture of purified water and glycerol and heated to 70°C. The remaining ingredients are heated together at 70°C. The two parts are added together and emulsified. The mixture is cooled and filled into containers.

e. Ointment

	weight (g)
active ingredient	12
white soft paraffin	88
	<u>100</u>

The white soft paraffin is melted at 60°C. The active ingredient is added and dispersed, allowed to cool, and filled into collapsible metal tubes.

f. Injection

	weight
active ingredient	250 mg
intralipid (Kabi-Vitrium)	10 ml

This composition is for example useful for intravenous injection.

g. Tablet

	mg/tablet
active ingredient	100
lactose	235
starch	50
polyvinylpyrrolidone	50
magnesium stearate	5
	<u>500</u>

Mix the active compound with the lactose and starch and wet granulate with a solution of the polyvinylpyrrolidone. Dry, sift, blend the granules with magnesium stearate and compress.

These formulations g), h) and i) are examples of those where the lipid compatibility of the derivatives of the invention leads to uptake in part direct into the lymph system with the advantage of avoiding direct passage to the liver already referred to.

h. Capsule (I)

	mg/capsule
active compound	200
lactose	189
sodium starch glycollate	8
polyvinylpyrrolidone	6
magnesium stearate	2
	<u>400</u>

Mix the active compound with the lactose and sodium starch glycollate and wet granule with a solution of the polyvinylpyrrolidone. Dry, sift, blend the granulate with the magnesium stearate and fill into hard gelatin capsules.



## i. Capsule (II)

	mg/capsule
active ingredient	100
arachis oil	500 ml

The active compound is dissolved in arachis oil and filled into soft or hard gelatin capsules.  
Arachis oil compositions are for example useful against ocular herpes infection.

## j. Topical preparation (liquid)

active ingredient	0.5%
propylene glycol	50.0%
ethanol	49.5%

The following are specific examples employing the above formulations:-

Example (1) Tablets containing 300 mg of LA-AZT. 1 or 2 tablets to be taken 4 to 6 hourly against herpes simplex or HIV and related retroviral infections.

Example (2) Hard gelatin capsules containing 150 mg of LA-AZT. 1 or 2 capsules to be taken 4 to 6 hourly against herpes simplex or retroviral infections.

Example (3) Syrup for oral administration containing 150 mg/ 5 ml LA-AZT. 5 ml to be taken 4 to 6 hourly against herpes simplex or retroviral infections.

Example (4) Solution for oral administration or intramuscular or intravenous injection containing 200 mg of LA-AZT in 3 ml, to be given 4 to 6 hourly against herpes simplex or retroviral infections. Intravenous injection for example is a preferred route for achieving therapeutic concentration in brain tissue across the blood brain barrier.

Example (5) Ointment or cream for topical administration against herpes simplex, containing 2% by weight LA-AZT. It may be applied to a human afflicted with a herpetic lesion at the afflicted sites at a rate of 1 to 5 mg/cm<sup>2</sup> three times daily for a period of one week. Reduction in the size of the lesion is noted after three days.

Examples (6) to (10) correspondingly, LA-AHT, is used individually in amounts and formulations as above. Further alternatives are corresponding preparations based on the ddC, ddA and D4T derivatives of synthesis examples 6 and 10 to 15 namely AK15 and NL1 to 6, for the same formulations and uses.

Example (11) Hard gelatin capsules containing 800 mg of LA-AZT. 1 or 2 capsules to be taken 4 to 6 hourly against herpes zoster.

Example (12) Tablets containing 300 mg of di-LA-acyclovir or o-LA-acyclovir (synthesis Examples (4) and (3); 1 or 2 tablets to be taken 4 to 6 hourly against virus infections. The n-LA-acyclovir of synthesis Example 5 is an alternative.

Example (13) Hard gelatin capsules containing 150 mg of the o-LA-acyclovir or other derivatives; 1 or 2 capsules to be taken 4 to 6 hourly against virus infections.

Example (14) Syrup for oral administration containing 150 mg/ 5 ml of the o-LA-acyclovir or other derivatives; 5 ml to be taken 4 to 6 hourly against virus infections.

Example (15) Ointment or cream for topical administration against virus infections containing 2% by weight of the o-LA-acyclovir or other derivatives.

Examples (16) to (18) Correspondingly, DHPG (gancyclovir) derivatives are used as set out in Examples (13) to (15).

Among the virus infections for which the above formulations (12) to (18) may be used are herpes simplex and herpes zoster.

## PHYSICAL PROPERTIES

Physical properties of certain of the new compounds have been measured as follows:-

TABLE 3

Partition co-efficients, water/chloroform	
Compound	% in chloroform phase
AK 2 LA - AZT	> 99%
AK 4 LA - AHT	> 99%
AK 6 LA - ACV	> 99%
AK 7 AZT (comparative)	15%
AK 8 AHT (comparative)	4%
AK 9 ACV (comparative)	< 1%
AK 11 di-LA-ACV	> 99%

TABLE 4

Absolute Solubilities		
	In Water	In CHCl <sub>3</sub>
AK 6 LA - ACV	0.014 g/l	> 50 g/l
AK 9 ACV	1.590 g/l	0.006 g/l

\* 98 to 99% pure, with 1% balance ethanol to stabilise.

As already discussed, the greatly enhanced solubility in chloroform of the fatty acid-ACV compounds compared to ACV itself, with comparable Figures shown by AZT/AHT, is important to the drug pharmacokinetics. Lipophilic compounds pass through cell membranes to act within the cytoplasm. Normally, water soluble compounds do not readily pass through cell membranes and require specific membrane transport mechanisms to achieve high intra-cellular concentrations. Further, the oil like properties of the fatty acid nucleoside analogue derivatives itself assists manufacture, formulation, and application (especially topically) for anti-viral formulations.

## TESTING FOR EFFICACY

### A. Inhibition of Virus Replication, Method

1. Mono-layers of baby hamster kidney cells (BHK-21) are prepared in Petri dishes containing ETC medium (Eagle's minimal essential medium + 10% tryptose phosphate broth + 10% new born calf serum).
2. After incubation the medium is poured off to leave the layer of cells attached to the Petri dish. A solution containing virus, specifically herpes simplex virus Type 1 (HSV 1) for example, in ETC medium is then added to the dish and incubated with intermittent rocking for one hour to allow the virus to adsorb to the cells of the mono-layer. Unadsorbed virus is then removed by pouring off the virus-containing medium and washing twice with new sterile medium. Then new medium is added to each Petri dish. At this stage various concentrations of drug can be added to the medium.
3. After a further 24 hour incubation to allow the virus to multiply in the cells to which it has become adsorbed, the cultures are treated to enable the amount of virus produced to be assayed. First the dishes are frozen to -70°C and thawed to disrupt the mono-layer and break up the cells. Then the medium and the disrupted cells are transferred to bijoux bottles and sonicated to break down the cells completely and to release all the virus particles the cells contain. The sonicated medium is then diluted with fresh sterile ETC medium to give concentrations of the original Petri dish fluid of  $10^{-2}$ ,  $10^{-4}$ ,  $10^{-6}$  and  $10^{-8}$ .

4. The number of virus particles (plaque forming units) in the dilutions is then assayed. A standard number of BHK cells (usually  $8 \times 10^6$ ) is added to 2 ml of the diluted solutions and incubated for one hour, with cautious, gentle shaking, and 8 ml of Eagle's minimal essential medium + 2% new born calf serum + carboxymethyl cellulose is then added to the cell suspension. This medium has gelling properties which prevent spreading of the virus through the medium and which encourage the formation of discrete plaques in cell mono-layers. The medium from each dilution is then dispensed into Petri dishes and incubated for two to three days.

5. The number of plaques formed is then counted and the amount of active virus expressed as the number of plaque forming units (pfus) as follows. The medium is poured off and the cell mono-layer is fixed using 10% formol saline and stained with dilute carbol fuchsin. The number of plaques in each Petri dish is counted. Based on the dilutions, the number of plaque forming virus particles in the original solutions is calculated.

#### B. Inactivation of Virus, Method

1. A virus solution containing  $2 \times 10^5$  pfu/ml is prepared in phosphate buffered saline (Dulbecco's A).
2. The compound to be tested is made up in Dulbecco's medium A in a suitable range of dilutions.
3. Equal volumes of the virus and drug solution are mixed together. Control solutions without drug are also used.
4. The virus and drug are incubated for an appropriate period of time, in this example, thirty minutes.
5. The numbers of active viral particles in the solution are assayed as in items 4 and 5 under A.

#### C. Results, AZT and AHT

AZT alone is known to have no effect as regards either the replication or the inactivation of herpes simplex virus, see for example, E. de Klerk et al, Biochemical Pharmacology 29: 1849, (1980) and the same is true of AHT. The inventors have, however, investigated the effects of LA and of LA-AZT and LA-AHT prepared as described herein, on herpes simplex, Type 1, Troisbel strain.

A first investigation was an assay to test whether LA-AZT or LA could inhibit viral replication and was set up using the method above. LA was found to have no effect at all. It was found that in complete contrast to either AZT alone or LA alone, the compound of LA-AZT was able to inhibit viral replication effectively. The new compound thus revealed properties possessed by neither of its constituents, inhibition of herpes virus replication. Like results are obtained when the AHT derivative is tested.

A second investigation was an assay to test whether LA-AZT and LA inactivated the herpes virus.

It was found that both LA and LA-AZT are able to inactivate the herpes simplex virus but the LA-AZT was more effective. The LA was active at concentrations 5-25  $\mu\text{g/ml}$  and LA-AZT at 1 to 3  $\mu\text{g/ml}$ , with corresponding results for AHT derivatives. Other virucidal fatty acids may, it follows, be expected to retain their activity when used as derivatives.

In summary, AZT and AHT alone have no effect on either inactivation or replication of simplex herpes I virus, and while LA alone has some effect in inactivating the herpes virus they are completely unable to inhibit viral replication. In contrast, compounds such as LA-AZT and the corresponding AHT derivatives are able both to inactivate the herpes virus and to inhibit its replication. The latter effect, in particular, could not have been predicted on the basis of the known separate effects of the two agents.

#### D. Results, ACV and materials generally

Using the above methods the results summarised in the following Table have been obtained in relation to ACV and derivatives thereof and also in relation to AZT and AHT.

TABLE 5

ANTIVIRAL ACTIVITY OF COMPOUNDS AGAINST HERPES SIMPLEX VIRUS I		
COMPOUND	INACTIVATION OF VIRUS (log <sub>10</sub> reduction in virus titre)	INHIBITION OF VIRAL REPLICATION (log <sub>10</sub> reduction in virus titre)
AK 2	2.2 @ 1 mg/l	no inhibition up to 100 mg/l
LA-AZT	2.8 @ 5 mg/l	
AK 4	2.75 @ 1 mg/l	no inhibition up to 100 mg/l
LA-AHT	2.9 @ 5 mg/l	
AK 6	no inactivation up to 50 mg/l	2.4 @ 1 mg/l
LA-ACV		4.2 @ 5 mg/l
AK 7	no inactivation up to 50 mg/l	no inhibition up to 100 mg/l
AZT		
AK 8	no inactivation at 10 mg/l 0.7 @ 25 mg/l	no inhibition of replication
AHT		
AK 9	no inactivation up to 50 mg/l	2 @ 0.1 mg/l
ACV		2.3 @ 0.5 mg/l
AK 11	0.15 @ 1 mg/l 1.55 @ 5 mg/l	1.69 @ 1 mg/l
di-LA-ACV		2.46 @ 5 mg/l
AK 13	1.3 @ 1 mg/l 1.8 @ 5 mg/l	2.45 @ 0.1 mg/l
n-LA-ACV		4.3 @ 1 mg/l

#### Anti-HIV activity assay

Results in relation to HIV virus were obtained by the method of Pauwels et al, Journal of Virological Methods 20, 309 to 321 (1988), as follows:

#### Method

MT-4 cells ( $2.5 \times 10^4$ /well) were infected with HIV (100 CCID<sub>50</sub>) and incubated in the presence of varying concentrations of the test compounds (added immediately after virus infection). After five days incubation at 37°C in a CO<sub>2</sub> incubator, the number of viable cells was assessed by the MTT (tetrazolium) method. Anti-viral activity and cytotoxicity of the compounds are expressed as ED<sub>50</sub> (50% effective dose) and CD<sub>50</sub> (50% cytotoxic dose), respectively. SI (selectivity index) = CD<sub>50</sub>/ED<sub>50</sub>.

Results

These are given in Table 6 and Table 7 below in relation to HTLV-III<sub>B</sub> and LAV-2<sub>ROD</sub> viruses.

TABLE 6

Results: HTLV-III <sub>B</sub>			
	CD <sub>50</sub> (μg/ml)	ED <sub>50</sub> (μg/ml)	Selectivity index (SI)
AK 2	9.4929	0.000365	26008
AK 4	18.1641	> 40	< 1
AK 6	30.7272	> 40	< 1
AK 7	1.5368	0.000372	4131
AK 8	771.20	13.10	59
AK 9	300.0550	> 1000	< 1
AK 11	13.0923	> 40	< 1
AK 15	16.50	0.064	256
AK 16 (D4T)	44.41	0.077	574

TABLE 7

Results: LAV-2 <sub>ROD</sub>			
	CD <sub>50</sub> (μg/ml)	ED <sub>50</sub> (μg/ml)	Selectivity index (SI)
AK 2	5.9512	0.000744	7999
AK 4	17.4332	> 40	< 1
AK 6	62.4324	> 100	< 1
AK 7	1.5633	0.000534	2928
AK 8	330.40	21.80	15
AK 9	397.5805	> 10000	< 1
AK 11	10.1674	> 40	< 1
AK 15	24.23	0.064	378
AK 16 (D4T)	46.65	0.039	1196

**Claims**

1. Anti-virals wherein a linoleoyl or alpha-linolenyl acyl group or a fatty acyl group of the n-9 or n-7 series unsaturated fatty acids (C:16 upwards), is borne directly on a hydroxy or amino group of the sugar/sugar analogue or heterocyclic moiety of a nucleoside or nucleoside analogue.
2. Derivatives of pentose nucleosides, or of analogues thereof having an acyclic group in place of the pentose which acyclic group has a hydroxy substituted carbon at the position corresponding to the 5' position of the pentose, said derivatives being esters at the 5' position of the nucleoside or the hydroxy substituted carbon of the analogue as the case may be formed with fatty acids as set out in claim 1.
3. Fatty acyl AZT or AHT esters wherein linoleoyl or alpha-linolenyl or an unsaturated fatty acyl group of the n-9 or n-7 series fatty acids (C:16 upwards) is borne directly on the hydroxy group of the AZT or AHT.

4. Fatty acyl acyclovir esters, amides or ester/amides wherein linoleoyl or alpha-linolenyl or an unsaturated fatty acyl group of the n-9 or n-7 series fatty acids (C:16 upwards) is borne directly on the hydroxy and/or on the amino group of the acyclovir.
5. Fatty acyl ddC, ddA, D4T and gancyclovir derivatives wherein linoleoyl or alpha-linolenyl or an unsaturated fatty acyl group of the n-9 or n-7 series fatty acids (C:16 or upwards) is borne directly on the hydroxy and/or on an amino group of the ddC, ddA, D4T or gancyclovir.
6. Fatty acyl derivatives according to any of claims 3, 4 and 5 wherein the fatty acid of the acyl group is as defined in claim 1.
7. An anti-viral medicament comprising a fatty acyl derivative as claimed in any of claims 1 to 6 and optionally a pharmaceutically acceptable diluent or carrier.
8. Use of a fatty acyl derivative as claimed in any of claims 1 to 6 in the preparation of a medicament for use in the treatment of a viral infection.
9. Use of a fatty acyl derivative as claimed in any of claims 3 to 6 in the preparation of a medicament for use in the treatment of herpes.
10. Use as claimed in claim 9 in the preparation of a medicament for use in the treatment of herpes zoster or herpes simplex.
11. Use as claimed in claim 9 or claim 10 wherein said fatty acyl derivative is a derivative of acyclovir or gancyclovir.
12. Use as claimed in claim 8 with the purpose of enhancing transport of anti-virals across lipid barriers in the body, and especially from the gut into the lymph system, or into cells from the extra-cellular fluid, or across the blood brain barrier, or percutaneously in topical preparations, or of inhibiting glucuronidation or other metabolic modifications reducing drug efficacy.
13. Use of a fatty acyl derivative as claimed in any of claims 1, 2, 3 and 5 or claim 6 as dependent upon any of claims 1, 2, 3 and 5 in the preparation of a medicament for use in the treatment of viral infections with the purpose of enhancing transport of nucleoside analogue anti-virals across lipid barriers in the body, and especially from the gut into the lymph system or into cells from the extra-cellular fluid, or across the blood brain barrier, or percutaneously in topical preparations, or of inhibiting glucuronidation or other metabolic modifications reducing drug efficacy, without loss of replication inhibiting activity.

Claims for the following Contracting States : ES, GR

1. A method of making an anti-viral compound wherein linoleic acid (LA) or alpha-linolenic acid (ALA) or an unsaturated fatty acid of the n-9 or n-7 series fatty acids (C:16 upwards) is used to acylate a hydroxy or amino group of a sugar/sugar analogue or heterocyclic moiety of a nucleoside or nucleoside analogue.
2. A method of making a derivative of a pentose nucleoside or analogue thereof having an acyclic group in place of the pentose which acyclic group has a hydroxy-substituted-carbon at the position corresponding to the 5' position of the pentose, wherein the acyl group of an at least triply unsaturated fatty acid as defined in claim 1 is used to acylate the 5' hydroxy group or the corresponding hydroxy group of the analogue as the case may be.
3. A method of making an AZT or AHT ester wherein linoleic acid or alpha-linolenic acid or an unsaturated fatty acid of the n-9 or n-7 series fatty acids (C:16 upwards) is used to acylate the hydroxy group of the AZT or AHT.
4. A method of making an acyclovir ester, amide or ester/amide wherein linoleic acid or alpha-linolenic acid or an unsaturated fatty acid of the n-9 or n-7 series fatty acids (C:16 upwards) is used to acylate the hydroxy and/or the amino group of the acyclovir.
5. A method of making a ddC, ddA, D4T or gancyclovir derivative wherein linoleic acid or alpha-linolenic acid or an unsaturated fatty acid of the n-9 or n-7 series fatty acids (C:16 upwards) is used to acylate a hydroxy and/or amino group of the ddC, ddA, D4T or gancyclovir.

6. A method according to any of claims 3, 4 and 5 wherein the fatty acid of the acyl group is as defined in claim 1.
7. Use of a fatty acyl derivative obtainable by a method as claimed in any of claims 1 to 6 in the preparation of a medicament for use in the treatment of a viral infection.
8. Use of a fatty acyl derivative obtainable by a method as claimed in any of claims 3 to 6 in the preparation of a medicament for use in the treatment of herpes.
9. Use as claimed in claim 8 in the preparation of a medicament for use in the treatment of herpes zoster or herpes simplex.
10. Use as claimed in claim 7 with the purpose of enhancing transport of anti-virals across lipid barriers in the body, and especially from the gut into the lymph system, or into cells from the extra-cellular fluid, or across the blood brain barrier, or percutaneously in topical preparations, or of inhibiting glucuronidation or other metabolic modifications reducing drug efficacy.
11. Use of a fatty acyl derivative obtainable by a method as claimed in any of claims 1, 2, 3 and 5 or claim 6 as dependent upon any of claims 1, 2, 3 and 5 in the preparation of a medicament for use in the treatment of viral infections with the purpose of enhancing transport of nucleoside analogue anti-virals across lipid barriers in the body, and especially from the gut into the lymph system or into cells from the extra-cellular fluid, or across the blood brain barrier, or percutaneously in topical preparations, or of inhibiting glucuronidation or other metabolic modifications reducing drug efficacy, without loss of replication inhibiting activity.
12. Use as claimed in claim 8 or claim 9, wherein said fatty acyl derivative is a derivative of acyclovir or gancyclovir.



European Patent  
Office

# PARTIAL EUROPEAN SEARCH REPORT

Application Number

which under Rule 45 of the European Patent Convention EP 95 11 4526 shall be considered, for the purposes of subsequent proceedings, as the European search report

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. CL.5)
Y	EP-A-0 217 580 (THE WELLCOME FOUNDATION) * abstract * * claims 1-28 *	1-13	C07H19/06 C07D473/00 A61K31/70 A61K31/52
Y	EP-A-0 085 424 (SYNTEX U.S.A. INC.) * abstract *	1-13	
Y	EP-A-0 082 668 (BEECHAM GROUP PLC) * abstract *	1-13	
Y	EP-A-0 056 265 (DR. THILO & CO. GMBH) * abstract *	1-13	
A	CHEMICAL ABSTRACTS, vol. 105, no. 11, 15 September 1986, Columbus, Ohio, US; abstract no. 97876n, N. ITO ET AL. '5-Fluorouracil Derivatives as Pharmaceuticals.' page 678 ; * abstract * & JP-A-61 044 897 (TERUMO CORP.) 4 March 1986  -/-	1	
			TECHNICAL FIELDS SEARCHED (Int. CL.5)
			C07H C07D A61K
INCOMPLETE SEARCH			
<p>The Search Division considers that the present European patent application does not comply with the provisions of the European Patent Convention to such an extent that it is not possible to carry out a meaningful search into the state of the art on the basis of some of the claims</p> <p>Claims searched completely:</p> <p>Claims searched incompletely:</p> <p>Claims not searched:</p> <p>Reason for the limitation of the search:</p> <p>see sheet C</p>			
Place of search		Date of completion of the search	Examiner
THE HAGUE		14 November 1995	Scott, J
CATEGORY OF CITED DOCUMENTS			
<p>X : particularly relevant if taken alone</p> <p>Y : particularly relevant if combined with another document of the same category</p> <p>A : technological background</p> <p>O : non-written disclosure</p> <p>P : intermediate document</p> <p>T : theory or principle underlying the invention</p> <p>E : earlier patent document, but published on, or after the filing date</p> <p>D : document cited in the application</p> <p>L : document cited for other reasons</p> <p>&amp; : member of the same patent family, corresponding document</p>			

EPO FORM 1503 01.87 (P04.001)





# PARTIAL EUROPEAN SEARCH REPORT

Application Number

EP 95 11 4526

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int.CL5)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
A	CHEMICAL ABSTRACTS, vol. 98, no. 25, 20 June 1983, Columbus, Ohio, US; abstract no. 214136w, Y.NUNOKAWA ET AL. '1-beta-D-Ribofuranosyl-4,5-Substituted Imidazole' page 437 ; * abstract * & JP-A-58 013 394 (AJINOMOTO CO., INC.) 25 January 1983 -----	1	
A	CHEMICAL ABSTRACTS, vol. 76, no. 7, 14 February 1972, Columbus, Ohio, US; abstract no. 24533k, M.HONJO ET AL. 'N4,2',3',5'-Tetraalkylanoylcytidines' page 345 ; * abstract * & JP-A-71 037 827 (TAKEDA CHEMICAL INDUSTRIES LTD.) 8 November 1971 -----	1	TECHNICAL FIELDS SEARCHED (Int.CL5)



EP 95 11 4526

-C-

Remark: Although claims 12 and 13  
are directed to a method of  
treatment of the human/animal  
body (Art. 52(4) EPC) the search  
has been carried out and based on  
the alleged effects of the  
compound/composition